
## Clostridium Difficile (C-Diff) Ordering for Diagnosis


Clostridium difficile associated disease (CDAD) continues to be a significant problem within the community and healthcare institutions. A systematic and evidence-based approach to diagnosis of CDAD can help minimize unnecessary testing and still provide the highest yield in diagnosis. To implement an appropriate approach to PCR (or other target amplification testing), incorporates Millennium Discern Rules approved by the Care Management Team and conforms to Infection Prevention Policies. They reflect current standard of care and assure appropriate test utilization. NOTE: PCR testing may be performed only on diarrheal stool. Any formed samples received by the laboratory will be rejected for testing.

## laboratory diagnosis of CDAD, the Infectious Diseases Division of the Clinical Laboratories of Banner Health has initiated a monitoring and control system for orders for C. difficile Toxin testing. The system includes a switch to **PHYSICIAN POSITIVE TEST and subsequent testing** $\bigcirc$ CLOSTRIDIUM DIFFICILE DX Allowance of only 1 positive PCR test per patient per episode You are placing an order for a C. diff Toxin (stool) study and the patient has had a (within 14 days). Subsequent requests for positive result within the last 14 days C difficile toxin testing during this time frame will be cancelled Allowance of only 1 positive test per patient per episode. Additional orders for toxin testing will be cancelled automatically automatically. A single positive test confirms CDAD when the right symptoms are present. Ordering toxin testing on stools as a test-of-cure is unnecessary and contraindicated - including for the purpose of terminating Contact Isolation (see Infection Prevention Policies). A single positive test confirms Clostridium difficile associated disease (CDAD) when symptoms are present. Ordering testing on stools as a test-of-cure is unnecessary and contraindicated – including for the purpose of terminating Contact Isolation (see Infection Prevention Policies).

Additional information related to this alert may be found by clicking on the "Reference" button on the alert.



When an order has been placed but result has not been posted yet on an already active order.




## Clostridium Difficile (C-Diff) Ordering for Diagnosis

## **NEGATIVE TEST and subsequent testing**

If one test by PCR (or other target amplification method) has been ordered on one encounter and results are negative, subsequent testing will not be allowed for 7 days from the most recent result.

Due to sensitivity of these methods, no further testing is required. All requests for more than 1 test will be cancelled automatically until 7 days has passed from the most recent result.

Additional information related to this alert may be found by clicking on the "Reference" button on the alert.



Other information: if the laboratory has cancelled a formed stool specimen, the Cancel Reason will show as:

Spec unacceptable for testing

And in PowerChart, this additional information will be seen as well:

